


CLAIMS

1. A compound of general formula (I) below:




- 5 wherein
- each R group is, at one or more of the positions 4, 5, 6 and 7 of the indolinone ring and independently from each other, a straight or branched C₁-C₄ alkyl or alkoxy group or a halogen atom;
- 10 each R₁ group is, the same or different and at one or more of the positions of the pyrrole ring, a C₁-C₄ alkyl or a group of general formula -(CH₂)_p-CO₂R', -(CH₂)_p-CONR'R" or -CONH-(CH₂)_p-CONR'R" wherein p is 0, 1, 2 or 3, the alkylene -(CH₂)_p- chain is optionally substituted by hydroxy, and R' and R" are selected, each independently, from hydrogen or
- 15 straight or branched C₁-C₄ alkyl optionally substituted by hydroxy or, taken together with the nitrogen atom to which they are attached, R' and R" may form a pyrrolidino, piperidino or morpholino group;
- 20 m is 0 or an integer from 1 to 4;
n is 0 or an integer from 1 to 3;
or pharmaceutically acceptable salts thereof.
2. A compound according to claim 1 wherein the pyrrole ring is substituted by one or more of the groups selected
- 25 from methyl, carboxy, ethoxycarbonyl, carboxyethyl, N,N-diethyl-aminocarbonyl, N-[(2-diethylamino)ethyl]carboxamide or N-[2-hydroxy-3-morpholin-4-ylpropyl]carboxamide.
3. A compound according to claim 1 which is 3-[(3,5-dimethyl-1H-pyrrol-2-yl) [¹⁴C]methylene-1,3-dihydro-2H-indol-
- 30 2-one; 5-[(1,2-dihydro-2-oxo-3H-indol-3-


- 25 -

ylidene) [¹⁴C]methyl]-2,4-dimethyl-1H-pyrrole-3-propionic acid; N-[(2-diethylamino)ethyl]-5-[(5-fluoro-1,2-dihydro-2-oxo-3H-indol-3-ylidene) [¹⁴C]methyl]-2,4-dimethyl-1H-pyrrole-3-carboxamide; 3-{5-methyl-2-[(Z)-(2-oxo-1,2-dihydro-3H-indol-3-ylidene) [¹⁴C]methyl]-1H-pyrrol-3-yl}propanoic acid; and 5-[(Z)-(5-fluoro-2-oxo-1,2-dihydro-3H-indol-3-ylidene) [¹⁴C]methyl]-N-[(2S)-2-hydroxy-3-morpholin-4-ylpropyl]-2,4-dimethyl-1H-pyrrole-3-carboxamide.


4. A process for preparing a compound of formula (I) according to claim 1 which process comprises:
- 10 a) reacting dimethyl- [¹⁴C]formamide with a suitable pyrrole derivative of formula (II), in the presence of diphosphoryl-chloride



- 15 wherein R_1 and n are as defined in claim 1, so as to obtain a compound of formula (III)



- and optionally converting a compound of formula (III) into another compound of formula (III);
- 20 b) reacting under basic conditions the compound of formula (III) with an oxindole derivative of formula (IV)




- wherein R and m are as defined in claim 1, so as to obtain a compound of formula (I) and, optionally converting it
- 25 into another compound of formula (I) and/or into a pharmaceutically acceptable salt thereof.


5. A process according to claim 4 wherein, in step (b), basic conditions are obtained by means of pyrrolidine.

6. A compound of formula (IIIa) or (IIIb) below


- 26 -



(IIIa)



(IIIb)

wherein R_1 is a hydrogen atom or a group selected from
 $-(CH_2)_2-CO_2H$, $-CO_2H$, $-CO_2CH_2CH_3$, $-CONH-(CH_2)_2-N(CH_2CH_3)_2$ and
 $-CONH-CH_2-CH(OH)-CH_2-N$ 

- 5 7. Use of a compound of formula (I), as defined in claim 1, for absorption, distribution, metabolism and excretion (ADME) studies.